

UK- 427,857 (maraviroc)



Drug Class: Entry and Fusion Inhibitors

Drug Description

UK-427,857 is a chemokine receptor antagonist that acts as an entry inhibitor. It is designed to prevent HIV infection of CD4 cells by blocking the CCR5 coreceptor from binding to HIV. [1]

HIV/AIDS-Related Uses

UK-427,857 is an investigational entry inhibitor that binds to a protein on the membrane of T cells called CCR5. The CCR5-tropic variant of the virus is common in earlier HIV infection, so UK-427,857 may be most useful in acute and early infection.[2]

Pharmacology

UK-427,857 binds to the CCR5 receptor, preventing HIV from binding to this receptor. When the CCR5 receptor is unavailable, CCR5-tropic HIV cannot engage a CD4 cell to infect the cell. The CCR5-tropic variant of the virus is common in earlier HIV infection, while viruses adapted to use the CXCR4 receptor gradually become dominant as HIV infection progresses. Thus, UK-427,857 may be most useful in acute and early infection, but further studies are needed.[3]

In a small Phase I study from 2003, 24 HIV infected adults with CCR5-tropic HIV were randomized to receive 25 mg UK-427,857 once daily, 100 mg twice daily, or placebo. Steady state drug levels were reached within 7 days, with more favorable drug levels achieved in the fasted state. By Day 14, those receiving 100 mg doses had experienced a viral load decline of more than 20-fold, compared to a nearly 3-fold reduction in the 25 mg group. The drug was well tolerated, and viral load did not rebound immediately upon cessation of the drug, indicating that a proportion of the receptors remain blocked for some time.[4]

A second placebo-controlled Phase I study of UK-427,857 demonstrated greater viral load declines with monotherapy than with placebo. Based on the results of this study, the researchers recommend a dose of 100 mg twice daily or higher be used. A similar study comparing a range of doses also showed viral load reductions of more

than 10-fold at doses above 100 mg once or twice daily, with no effect of food on the drug's antiviral efficacy.[5]

It is not known if the CCR5 protein on T cells or HIV can become resistant to UK-427,857 or how quickly such resistance could develop.[6] In an in vitro study using six primary CCR5 HIV-1 isolates, those able to replicate in the presence of high UK-427,857 concentrations emerged gradually after multiple passages of all isolates. Two isolates resistant to UK-427,857 continued to use the CCR5 receptor and one isolate developed the ability to use the CXCR4 receptor. In the viruses that remained R5-tropic, two different sets of mutations developed in the gp120 V3 loop region; this and other data suggest that changes in viral tropism are independent of UK-427,857.[7] [8] All CCR5 antagonists bind to CCR5 in a pocket formed by transmembrane helices and extracellular loop 2 (ECL2); it appears that subtle differences in occupation of the binding pocket may block replication of some HIV strains. As a result, scientists are optimistic that resistance to an HIV coreceptor antagonist will not necessarily lead to drug class resistance.[9]

Adverse Events/Toxicity

In 80 HIV infected adults, UK-427,857 was well tolerated.[10] In another Phase I study enrolling 54 HIV uninfected adults taking 100 or 300 mg UK-427,857 twice daily, no adverse effects were observed on white or red blood cell counts or any other blood tests.[11]

In a study of UK-427,857's effects on the QTc interval, single doses of UK-427,857 (at doses of 100, 300, or 900 mg) were given to healthy volunteers. Dizziness was reported when UK-427,857 was given at a 900 mg single dose; other side effects reported from this study include headache, postural hypotension, nausea, and cystitis.[12]

Drug and Food Interactions

In a small Phase I study, UK-427,857's antiviral effectiveness against HIV was unaffected by

UK- 427,857 (maraviroc)



Drug and Food Interactions (cont.)

food.[13]

In HIV infected adults, a dose of 150 mg UK-427,857 twice daily following a high fat meal was evaluated in order to assess the effect of food on antiviral efficacy. UK-427,857's plasma concentrations were similar to those seen in healthy volunteers under similar food restrictions.[14]

A four-cohort study, with administration of single doses of UK-427,857, was conducted in 29 HIV infected people who were stable for 3 months on one of four protocol-chosen antiretroviral regimens. Cohort 1 received efavirenz, abacavir sulfate, and lamivudine; cohort 2 received efavirenz, didanosine, and tenofovir disoproxil fumarate (tenofovir DF); cohort 3 received nevirapine, lamivudine, and tenofovir DF; and cohort 4 received lopinavir/ritonavir, stavudine, and lamivudine. Participants received a single oral dose of UK-427,857; serial blood draws were conducted over 12 hours post-dose. Results suggested that taking UK-427,857 with regimens containing efavirenz led to an approximately 50% reduction in UK-427,857. Conversely, UK-427,857 exposure doubled in the cohort receiving lopinavir/ritonavir. Additional dose adjustment studies are needed to determine optimal dosages for UK-427,857, especially when taken with other antiretrovirals.[15]

Clinical Trials

For information on clinical trials that involve UK-427,857 (maraviroc), visit the ClinicalTrials.gov web site at <http://www.clinicaltrials.gov>. In the Search box, enter: UK-427,857 (maraviroc) AND HIV Infections.

Dosing Information

Mode of Delivery: Oral.[16]

Dosage Form: UK-427,857 has been studied at doses of 25, 50, 100, 150, 300, and 900 mg.[17] [18] Once- and twice-daily administrations of UK-427,857 have been studied.[19]

Further Reading

Fatkenheuer G, Pozniak AL, Johnson M, Plettenberg A, Staszewski S, Hoepelman IM, Saag M, Goebel F, Rockstroh J, Dezube B, Jenkins TM, Medhurst C, Sullivan JF, Ridgway C, Abel S, Youle M, van der Ryst E. Evaluation of dosing frequency and food effect on viral load reduction during short-term monotherapy with UK-427,857, a novel CCR5 antagonist. 15th International AIDS Conference, Bangkok, Abstract TuPeB4489, 2004.

Marks K, Gulick RM. New Antiretroviral Agents for the Treatment of HIV Infection. *Curr Infect Dis Rep*. 2004 Aug;6(4):333-339. PMID: 15265463

Muirhead G, Pozniak A, Gazzard B, Nelson M, Moyle G, Ridgway C, Taylor-Worth R, Russell D. A Novel Probe Drug Interaction Study to Investigate the Effect of Selected ARV Combinations on the Pharmacokinetics of a Single Oral Dose of UK-427,857 in HIV Positive Subjects. 12th Conference on Retroviruses and Opportunistic Infections, Boston, Abstract 663, 2005.

Walker D, Abel S, Comby P, Muirhead G, Nedderman A, Smith DA. Species differences in the disposition of the CCR5 antagonist, UK-427,857, a new potential treatment for HIV. *Drug Metab Dispos*. 2005 Apr;33(4):587-95 PMID: 15650075

Westby M, Smith-Burchnell C, Mori J, et al. In vitro escape of R5 primary isolates from the CCR5 antagonist, UK-427,857, is difficult and involves continued use of the CCR5 receptor. *Antivir Ther*. 2004;9:S10.

Manufacturer Information

UK-427,857 (maraviroc)
Pfizer Inc
235 East 42nd Street
New York, NY 10017-5755
(800) 438-1985

For More Information

Contact your doctor or an AIDSinfo Health

UK- 427,857 (maraviroc)



For More Information (cont.)

Information Specialist:

- Via Phone: 1-800-448-0440 Monday - Friday, 12:00 p.m. (Noon) - 5:00 p.m. ET
- Via Live Help: http://aidsinfo.nih.gov/live_help Monday - Friday, 12:00 p.m. (Noon) - 4:00 p.m. ET

References

1. Intl HIV Drug Resistance Wkshp - 13th, 2004. Abstract 6.
2. Intl HIV Drug Resistance Wkshp - 13th, 2004. Abstract 6.
3. Intl HIV Drug Resistance Wkshp - 13th, 2004. Abstract 6.
4. Interscience Conf on Antimicrobial Agents and Chemotherapy - 43rd, 2003. Abstract H-443.
5. Intl AIDS Conf - 15th, 2004. Abstract TuPeB4479.
6. Interscience Conf on Antimicrobial Agents and Chemotherapy - 43rd, 2003. Abstract H-443.
7. Intl HIV Drug Resistance Wkshp - 13th, 2004. Abstract 6.
8. J Gen Virol. - 2000;81:2899-904
9. Conf Retroviruses Opportunistic Infect. - 12th, 2005. Abstract 96.
10. Intl AIDS Conf - 15th, 2004. Abstract TuPeB4489.
11. Interscience Conf on Antimicrobial Agents and Chemotherapy - 43rd, 2003. Abstract H-874.
12. Intl AIDS Conf - 15th, 2004. Abstract TuPeB4605.
13. Intl AIDS Conf - 15th, 2004. Abstract TuPeB4489.
14. Intl AIDS Conf - 15th, 2004. Abstract TuPeB4489.
15. Conf Retroviruses Opportunistic Infect. - 12th, 2005. Abstract 663.
16. Interscience Conf on Antimicrobial Agents and Chemotherapy - 43rd, 2003. Abstract H-874.
17. Intl AIDS Conf - 15th, 2004. Abstract TuPeB4605.
18. Intl AIDS Conf - 15th, 2004. Abstract TuPeB4489.
19. Interscience Conf on Antimicrobial Agents and Chemotherapy - 43rd, 2003. Abstract H-443.